

MCA SET I

Q1. $\tan 1^\circ \tan 2^\circ \tan 3^\circ \tan 4^\circ \dots \tan 89^\circ$ is equal to

- (A) 1
- (B) 0
- (C) ∞
- (D) $\frac{1}{2}$

Q2. The minimum value of $3 \cos x + 4 \sin x + 5$ is

- (A) 5
- (B) 9
- (C) 7
- (D) 0

Q3. The equation $\sin x \cos x = 2$ has

- (A) One solution
- (B) Two solutions
- (C) Infinite solutions
- (D) No solution

Q4. If the sides of the triangle be 6, 10, and 14 then the triangle is:

- (A) Obtuse angled
- (B) Acute angled
- (C) Right angled
- (D) Equilateral

Q5. In a ΔABC , $\sin A : \sin B : \sin C = 1 : 2 : 3$. If $b = 4$ cm, the perimeter of the triangle is

- (A) 6 cm
- (B) 24 cm
- (C) 12 cm
- (D) 8 cm

Q6. $1 + \cot^2(\sin^{-1}x)$ is equal to

- (A) $\frac{1}{2x}$
- (B) x^2
- (C) $\frac{1}{x^2}$

(D) $\frac{2}{x}$

Q7. Angle of elevation of the sun when the shadow of the pole is $\sqrt{3}$ times the height of the pole is

- (A) 60°
- (B) 30°
- (C) 45°
- (D) 15°

Q8. If $a \times b = b \times c \neq 0$ and $a + c \neq 0$, then

- (A) $(a + c) \perp b$
- (B) $(a + c) \parallel b$
- (C) $a + c = b$
- (D) None of the above

Q9. If the normals at two points P and Q of a parabola $y^2 = 4ax$ intersect at a third point R on the curve then the product of the ordinates of P and Q is

- (A) $4a^2$
- (B) $2a^2$
- (C) $-4a^2$
- (D) $8a^2$

Q10. If retardation produced by air resistance is one tenth of the acceleration due to gravity the time to return from maximum height

- (A) Decreases by 9%
- (B) Increases by 11%
- (C) Decreases by 11%
- (D) Increases by 9%

Q11. If the line $y = 2x + k$ is a tangent to the curve $x^2 = 4y$ then k is equal to

- (A) 4
- (B) $1/2$
- (C) -4
- (D) $-1/2$

Q12. The latus rectum of the hyperbola $9x^2 - 16y^2 - 18x - 32y - 151 = 0$ is

- (A) $9/4$
- (B) 9
- (C) $3/2$
- (D) $9/2$

Q13. The eccentricity of the ellipse $\frac{(x-1)^2}{9} + \frac{(y+1)^2}{25} = 1$ is

- (A) 4/5
- (B) 3/5
- (C) 5/4
- (D) Imaginary

Q14. The line $x \cos \alpha + y \sin \alpha = p$ will be a tangent to the conic $\frac{x^2}{a^2} + \frac{y^2}{b^2} = 1$ if

- (A) $p^2 = a^2 \sin^2 \alpha + b^2 \cos^2 \alpha$
- (B) $p^2 = a^2 + b^2$
- (C) $p^2 = b^2 \sin^2 \alpha + a^2 \cos^2 \alpha$
- (D) None of the above

Q15. Curve $xy = c^2$ is said to be

- (A) Parabola
- (B) Rectangular hyperbola
- (C) Hyperbola
- (D) Ellipse

Q16. If you want to kick a football to the maximum distance, the angle at which it should be kicked is (assuming no air resistance)

- (A) 45°
- (B) 90°
- (C) 30°
- (D) 60°

Q17. The area of the triangle formed by the lines joining the vertex of the parabola $x^2 = 12y$ to the ends of its latus rectum is

- (A) 12 sq. units
- (B) 16 sq. units
- (C) 18 sq. units
- (D) 24 sq. units

Q18. If the vectors $3i + \lambda j + k$ and $2i - j + 8k$ are perpendicular then λ is

- (A) -14
- (B) 7
- (C) 14
- (D) 1/7

Q19. The time taken for a projectile thrown with a velocity v cm/s at an angle α with the horizontal to attain the maximum height is given by

(A) $\frac{v}{g}$

(B) $\frac{v}{\sin \alpha}$

(C) $\frac{v \sin \alpha}{g}$

(D) $(v \sin \alpha) g$

Q20. In a city 20% of the population travel by car, 50% travel by bus and 10% travel by both car and bus. Then, persons travelling by car or bus is

- A. 80%
- B. 40%
- C. 60%
- D. 70%

Ans (C)

Q21. If $A = \{2, 4, 5\}$ and $B = \{7, 8, 9\}$ the $n(A \times B)$ is equal to

- (A) 6
- (B) 9
- (C) 3
- (D) 0

Q22. In an exam, 70% of students passed in maths, 80% passed in physics, 75% passed in chemistry, 85% passed in Biology, and $x\%$ passed in all the four subjects. The minimum value of x is:

- (A) 10
- (B) 12
- (C) 15
- (D) None of the above

Q23. If A , B and C are any three sets then $A - (B \cup C)$ is equal to :

- A. $(A - B) \cup (A - C)$
- B. $(A - B) \cap (A - C)$
- C. $(A - B) \cup C$
- D. $(A - B) \cap C$

Q24. If a , b , c are in AP, then $\frac{(a-c)^2}{(b^2-4ac)}$ is equal to:

- A. 1
- B. 2
- C. 3

D. 4

Q25. The GM of the numbers $3, 3^2, 3^3, \dots, 3^n$ is :

- A. $3^{2/n}$
- B. $3^{(n+1)/2}$
- C. $3^{n/2}$
- D. $3^{(n-1)/2}$

Q26. If $a^2 + ab^2 + 16c^2 = 2(3ab + 6bc + 4ac)$ where a, b, c are non zero numbers. Then a, b, c are said to be in:

- (A) AP
- (B) GP
- (C) HP
- (D) None of the above

Q27. If the arithmetic, geometric and harmonic means between two positive real numbers be A, G, and H, respectively then

- (A) $A^2 = GH$
- (B) $H^2 = AG$
- (C) $G = AH$
- (D) $G^2 = AH$

Q28. If $\log_a x, \log_b x, \log_c x$ be in HP then a, b, c are in :

- A. AP
- B. HP
- C. GP
- D. None of the above

Q29. The number of straight lines that can be formed by joining 20 points no three of which are in same straight line except 4 of them which are in the same line

- A. 183
- B. 186
- C. 197
- D. 185

Q30. Number of ways in which any four letters can be selected from the word CORGOO is :

- A. 15
- B. 11

- C. 7
- D. None of the above

Q31. The number of divisors of 9600 including 1 and 9600 are:

- A. 60
- B. 58
- C. 48
- D. 46

Q32. In how many ways can 5 keys be put in a ring?

- A. $\frac{1}{2}4!$
- B. $\frac{1}{2}5!$
- C. $4!$
- D. $5!$

Q33. Let 'X' be a family of sets and R be a relation on X defined by 'A is disjoint from B'. Then, R is :

- (A) Reflexive
- (B) Symmetric
- (C) Anti-symmetric
- (D) Transitive

Q34. R is a relation from {11,12,13} to {8,10,12} defined by $y=x-3$, then R^{-1} is:

- A. {(8,11), (10,13)}
- B. {(11,18), (13,10)}
- C. {(10,13), (8,11)}
- D. None of the above

Q35. Let R be a reflexive relation on a set A and I be the identity relation on A. then,

- A. $R \subset I$
- B. $I \subset R$
- C. $R = I$
- D. None of the above

Q36. Let S be the set of all real numbers. Then, the relation $R = \{(a,b): 1 + ab > 0\}$ on S is :

- A. Reflexive and symmetric but not transitive
- B. Reflexive and transitive but not symmetric
- C. Symmetric, transitive but not reflexive
- D. Reflexive, transitive and symmetric

Q37. Let R & S be two non-void relations on a set A. Which of the following statements is false.

- A. R & S are transitive $\Rightarrow R \cup S$ is transitive
- B. R & S are transitive $\Rightarrow R \cap S$ is transitive
- C. R & S are symmetric $\Rightarrow R \cup S$ is symmetric
- D. R & S are reflexive $\Rightarrow R \cap S$ is reflexive

Q38. Let a relation R be defined by $R = \{ (4,5), (1,4), (4,6), (7,6), (3,7) \}$ then $R^{-1} \circ R$ is :

- A. $\{ (1,1), (4,4), (4,7), (7,4), (7,7), (3,3) \}$
- B. $\{ (1,1), (4,4), (7,7), (3,3) \}$
- C. $\{ (1,5), (1,6), (3,6) \}$
- D. None of the above

Q39. Function $f : R \rightarrow R, f(x) = x^2 + x$ is :

- A. One-one onto
- B. One-one into
- C. Many-one onto
- D. Many-one into

Q40. Domain of $f(x) = \log |\log x|$ is:

- A. $(0, \infty)$
- B. $(1, \infty)$
- C. $(0, 1) \cup (1, \infty)$
- D. $(-\infty, 1)$

Q41. The domain of definition of the function $y(x)$ given by $2^x + 2^y = 2$ is :

- A. $(0,1]$
- B. $[0,1]$
- C. $(-\infty, 0]$
- D. $(-\infty, 1)$

Q42. Which of the following function is invertible

- (A) $f(x) = 2^x$
- (B) $f(x) = x^3 - x$
- (C) $f(x) = x^2$
- (D) None of the above

Q43. Let $f : (2,3) \rightarrow (0,1)$ be defined by $f(x) = x - [x]$, then $f^{-1}(x)$ equals

- (A) $x - 2$
- (B) $x + 1$

- (C) $x - 1$
- (D) $x + 2$

Q44. If $a + b + c = 0$, $a \neq 0$, $a, b, c \in \mathbb{Q}$, then both the roots of the equation $ax^2 + bx + c = 0$ are:

- A. rational
- B. non-real
- C. irrational
- D. zero

Q45. A real root of an equation $\log_4 \{ \log_2 (\sqrt{(x+8)} - \sqrt{x}) \} = 0$ is :

- A. 1
- B. 2
- C. 3
- D. 4

Q46. For equation $3x^2 + px + 3 = 0$, $p > 0$, if one of the roots is square of the other, then p is equal to

- A. $\frac{1}{3}$
- B. 1
- C. 3
- D. $\frac{2}{3}$

Q47. The equation of motion of a vehicle is $s = t^2 - 2t$, where 't' is measured in hours and 's' in kilometres. When the distance travelled by the vehicle is 15 km, the velocity of the vehicle is :

- A. 2 km/h
- B. 4 km/h
- C. 6 km/h
- D. 8 km/h

Q48. The maximum value of $\left(\frac{1}{x}\right)^x$ is :

- A. $(e)^e$
- B. $(e)^{\frac{1}{e}}$
- C. $(e)^{-e}$
- D. $\left(\frac{1}{e}\right)^e$

Q49. If $f(x) = x + \frac{1}{x}$, $x > 0$, then its greatest value is :

- A. -2
- B. 0

- C. 3
- D. None of the above

Q50. The sum of coefficients in the expansion of $(x + 2y + 3z)^8$ is :

- A. 3^8
- B. 5^8
- C. 6^8
- D. 7^8

Q51. If the coefficients of r^{th} term and $(r + 4)^{th}$ term are equal in the expansion of $(1 + x)^{20}$, then the value of 'r' will be

- A. 7
- B. 8
- C. 9
- D. 10

Q52. If $n \in \mathbb{N}$ then $x^{2n-1} + y^{2n-1}$ is divisible by

- A. $x + y$
- B. $x - y$
- C. $x^2 + y^2$
- D. $x^2 + xy$

Q53. $\lim_{x \rightarrow 0} \left(\frac{x(e^x - 1)}{1 - \cos x} \right)$ is equal to

- A. 0
- B. ∞
- C. -2
- D. 2

Q54. If $\lim_{x \rightarrow 0} kx \operatorname{cosec} x = \lim_{x \rightarrow 0} x \operatorname{cosec} kx$ then 'k' is equal to

- A. 1
- B. -1
- C. ± 1
- D. ± 2

Q55. The function $y = e^{-|x|}$ is

- A. Continuous and differentiable at $x = 0$
- B. Neither continuous nor differentiable at $x = 0$
- C. Continuous but not differentiable at $x = 0$
- D. Not continuous but differentiable at $x = 0$

Q56. If $A = \begin{bmatrix} 1 & 1 \\ 0 & 1 \end{bmatrix}$ then A^n is equal to :

- A. $\begin{bmatrix} 1 & n \\ 0 & 1 \end{bmatrix}$
- B. $\begin{bmatrix} n & n \\ 0 & n \end{bmatrix}$
- C. $\begin{bmatrix} n & 1 \\ 0 & n \end{bmatrix}$
- D. $\begin{bmatrix} 1 & 1 \\ 0 & n \end{bmatrix}$

Q57. If A and B are square matrices of the same order then,

- A. $(AB)' = A' B'$
- B. $(AB)' = B' A'$
- C. $AB = 0$; if $|A| = 0$ or $|B| = 0$
- D. $AB = 0$; if $A = 1$ or $B = 1$

Q58. The derivative of $\sin^{-1}\left(\frac{2x}{1+x^2}\right)$ with respect to $\cos^{-1}\left(\frac{1-x^2}{1+x^2}\right)$ is

- A. -1
- B. 1
- C. 2
- D. 4

Q59. the derivative of $f(x) = x|x|$ is

- A. $2x$
- B. $-2x$
- C. $2x^2$
- D. $2|x|$

Q60. $\int \frac{dx}{e^x + e^{-x}}$ is equal to

- A. $\tan^{-1}(e^{-x}) + C$
- B. $\tan^{-1}(e^x) + C$
- C. $\log(e^x - e^{-x}) + C$
- D. $\log(e^{x^2} - e^{-x}) + C$

Q61. Value of $\int_0^{\frac{\pi}{2}} \frac{\sin x}{1 + \cos^2 x} dx$ is

- A. $\frac{\pi}{2}$
- B. $\frac{\pi}{4}$
- C. $\frac{\pi}{3}$

D. $\frac{\pi}{6}$

Q62. Area bounded by the curve $y = x^3$, 'x' axis and ordinates $x = 1$ and $x = 4$ is

A. $64 m^2$

B. $27 m^2$

C. $\frac{127}{4} m^2$

D. $\frac{255}{4} m^2$

Q63. The order of the differential equation of the family of all the concentric circles centered at (h,k) is

A. 1

B. 2

C. 3

D. 4

Q64. The slope of a curve at any point is the reciprocal of twice the ordinate at the point and it passes through the point (4,3). The equation of curve is

A. $x^2 = y + 5$

B. $y^2 = x - 5$

C. $y^2 = x + 5$

D. $x^2 = y - 5$

Q65. The area of the triangle formed by the lines

$$7x - 2y + 10 = 0, 7x + 2y - 10 = 0, y + 2 = 0$$

A. 8 sq. units

B. 12 sq. units

C. 14 sq. units

D. None of these

Q66. The equation $(x + y)^2 = (x^2 + y^2) = 0$ represents

A. Circle

B. Two lines

C. Two parallel lines

D. Two mutually perpendicular lines

Q67. If the sum of the slopes of the lines represented by the equation $x^2 - 2xy \tan A - y^2 = 0$ be 4, then $\angle A$ is equal to :

A. 0 degree

B. 45 degree

- C. 60 degree
- D. $\tan^{-1}(-2)$

Q68. Which the following is false ?

- (A) $x^2(1+x) > 0 \Leftrightarrow x > -1$ and $x \neq 0$
- (B) $x^3 + y^3 = 0 \Leftrightarrow x = y = 0$
- (C) $x = 3$ and $y = 5 \Rightarrow 2x + 4y = 26$
- (D) $x = \sqrt{16} \Rightarrow x^2 = 16$

Q69. Consider the statement : 'If it rains, the wind is blowing'. Which of the following statements does not express the same (with R as the statement that it is raining, and W as the statement that the wind is blowing, $R \Rightarrow W$) ?

- (A) A sufficient condition for rain is that the wind is blowing
- (B) A sufficient condition for the wind to blow is that it is raining
- (C) A necessary condition for rain is that the wind is blowing
- (D) If the wind is blowing, there will be no rain

Q70. In a group of 100 students, 25 study economics, 30 study political science and 5 study both subjects. How many students study neither economics nor political science?

- (A) 45
- (B) Unable to tell
- (C) 55
- (D) 50

Q71. Given the sets $A = \{2,3,4,5\}$, $B = \{1,2,3,4,7\}$ and $C = \{1,3,6,7\}$, which of the following statements is false?

- (A) $(A \setminus B) \cap C = \{2\}$
- (B) $A \cap C \subset B$
- (C) $(A \cup B) \cap C = \{1,3,7\}$
- (D) $2 \in A \cap B$

Q72. If $A = \begin{pmatrix} 0 & 1 \\ 2 & 3 \end{pmatrix}$ and $B = \begin{pmatrix} 1 & -1 \\ 5 & 2 \end{pmatrix}$, which of the following is false?

- A. $A + B = \begin{pmatrix} 1 & 0 \\ 7 & 5 \end{pmatrix}$
 B. $A^2 = \begin{pmatrix} 0 & 1 \\ 4 & 9 \end{pmatrix}$
 C. $AB = \begin{pmatrix} 5 & 2 \\ 17 & 4 \end{pmatrix}$
 D. $3A - 4B = \begin{pmatrix} -4 & 7 \\ -14 & 1 \end{pmatrix}$

Q73. If A, B and C are $n \times n$ matrices, which of the following equalities is invalid? Note: D' is the transpose of D

- (A) $(ABC)' = C' B' A'$
 (B) $(A + A)' = 2A'$
 (C) $((AB)^2)' = (B')^2 (A')^2$
 (D) $(A + A + 2B)' = 2B' + 2A'$

Q74. Which of the following statements is correct?

- A. A linear system with more equations than unknowns cannot have solutions
 B. It is possible to construct a linear system with exactly 5 different solutions
 C. Suppose **A** is $n \times n$, **x** is $n \times 1$, and $\mathbf{Ax} = 0$ has only the trivial solution. Then $\mathbf{Ax} = \mathbf{b}$ has solutions for any $n \times 1$, vector **b**
 D. A linear system can only have an infinite number of solutions if there are more variables than equations

Q75. For which values of t does the following linear equation system have infinitely many solutions

$$\begin{aligned} tx + y &= 1 \\ 6x + (t+1)y &= 3 \end{aligned}$$

- A. $t = -3$
 B. $t = 2$
 C. $t = 2$ and $t = -3$
 D. The system does not have infinitely many solutions for any value of t

Q76. Using Gaussian eliminations, the solutions of: $x + y + z = c$, $x + 2y + az = 2c$ and $x + 2y + bz = 2$ can be deduced from the augmented matrix

$$\begin{pmatrix} 1 & 1 & 1 & c \\ 0 & 1 & a-1 & c \\ 0 & 0 & b-a & 2(1-c) \end{pmatrix}$$

For which values of a , b , and c are there infinitely many solutions?

- (A) If $a \neq b$
- (B) If $a = b$ and $c = 1$
- (C) If $c = 1$
- (D) Never

Q77. The straight line in \mathbb{R}^3 through the point $(-1, 3, 3)$ pointing in the direction of the vector $(1, 2, 3)$ hits the $x_1 x_2$ - plane at the point:

- A. $(1, 3, 0)$
- B. Never
- C. $(2, -1, 0)$
- D. $(-2, 1, 0)$

Q78. Let L denote the line passing through the points $(0, 5)$ and $(4, 3)$. Which of the following points also lies on L ?

- A. $(8, 0)$
- B. $(12, -1)$
- C. $(11, 0)$
- D. $(-4, 6)$

Q79. Which of the following formulas is false (x , y , and z are positive)?

- (A) $(\ln x)^4 = 4 \ln x$
- (B) $\ln[(x + y)^{1/5} z^{2/3}]^{15} = 3 \ln(x + y) + \ln z$
- (C) $\ln x^5 - \ln x^3 = 2 \ln x$
- (D) $2 \ln \frac{x}{y} + \ln \frac{y^2}{x^2} = 0$

Q80. If $f(x) = \ln x$, $x > 0$, and $g(x) = 4 - x^2$, $x \in \mathbb{R}$ what is the range of $f(g(x))$?

- (A) $(-\infty, \ln 4)$
- (B) $(-\infty, 0)$
- (C) $(0, \infty)$
- (D) $(0, \ln 4)$

Q81. $\frac{(3^{100}+3^{98})}{(3^{100}-3^{98})}$ is equal to

- A. 3^{196}
- B. 99
- C. $\frac{3+3}{3-3}$
- D. $5/4$

Q82. If $x = a - b$ makes $x^2 - 2ax + m$ equal to 0, then m is

- A. $a + b$
- B. $a^2 - b^2$
- C. $a^2 + b^2$
- D. $a - b$

Q83. Which of the following factorizations is incorrect?

- A. $2a^2 - 5ab - 3b^2 = (2a + b)(a - 3b)$
- B. $x^6 - y^6 = (x^3 + y^3)(x^3 - y^3)$
- C. $25a^2 + 1 = (5a + 1)(5a - 1)$
- D. $a - 2\sqrt{ab} + b = (\sqrt{a} - \sqrt{b})^2$

Q84. If $p \in (0,1)$, then:

- A. $p > 1/p$
- B. $p^3 > p^2$
- C. $p > \sqrt{p}$
- D. $1/p > \sqrt{p}$

Q85. Which of the following statements is incorrect?

- (A) $\sqrt{a^2} = |a|$
- (B) $|a + b| \leq |a| + |b|$
- (C) $|a - b| \leq |a| - |b|$
- (D) $|a| - |b| \leq |a + b|$

Q86. In a sports league where no drawn games are possible, a team had 10 more wins than twice its losses. It played a total of 52 matches. How many did it lose?

- (A) 12
- (B) 10
- (C) 14
- (D) 16

Q87. The solution (s) of the equation $\frac{x^2-3x-10}{\sqrt{x-5}} = 0$ is / are :

- (A) Only $x = 5$
- (B) No solution
- (C) Only $x = -2$
- (D) $x = -2$ and $x = 5$

Q88. Solving $\frac{1}{p} + \frac{1}{q} = \frac{1}{T}$ for 'q' you get :

- (A) $\frac{pT}{p+T}$
- (B) $T - p$
- (C) $\frac{pT}{p-T}$
- (D) $\frac{1}{T} - \frac{1}{p}$

Q89. If $z = F(x, y) = x^2 - y^3$ and $x = t^2, y = 1 - t$, then $\left\{\frac{dz}{dt}\right\}_{t=0}$ is :

- (A) 3
- (B) 0
- (C) -3
- (D) 2

Q90. The function $z = xy^2 - y^3 + 2x^2y$ satisfies the equation $xz'x + yz'y = kz$ for $k =$

- (A) 2
- (B) 3
- (C) 4
- (D) For no value of k

Q91. Which of the following statements about systems of equations is correct?

- (A) Three equations with two unknowns never have a solution.
- (B) Three linear equations with three unknowns never have exactly two solutions.
- (C) Three linear equations with three unknowns always have a unique solution
- (D) Two equations with three unknowns always have a solution

Q92. Ogives for more than type and less than type distributions intersect at:

- (A) mean
- (B) median
- (C) mode
- (D) origin

Q93. If A and A^C are complementary events in a sample space S, then :

- (A) $P(A) + P(A^C) = 0$
- (B) $P(A) - P(A^C) = 0$
- (C) $P(A) + P(A^C) = 1$
- (D) $P(A) - P(A^C) = 1$

Q94. If mean of n observations is 'a'. If one observation 'b' is added, mean continues to remain 'a', then the value of 'b' is:

- (A) 0
- (B) 1
- (C) n
- (D) a

Q95. Let there be two data sets I and II of size 80 and 20 respectively. The combined arithmetic mean of the two data sets is 500. If the arithmetic mean of the data set I is 520, then the arithmetic mean of data set II is :

- (A) 480
- (B) 490

(C) 450

(D) 420

Q96. The mean of 50 observation is 40 and standard deviation (s.d.) 8. If 4 is added to each observation, then the new mean and standard deviation (s.d.) are :

(A) mean = 40, s.d. = 8

(B) mean = 44, s.d. = 12

(C) mean = 44, s.d. = 8

(D) mean = 40, s.d. = 12

Q97. Let $f(x) = \int_1^x \sqrt{2 - t^2} dt$ then real roots of the equation $x^2 - f'(x) = 0$ are

A. ± 1

B. $\pm \frac{1}{\sqrt{2}}$

C. $\pm \frac{1}{2}$

D. 0 and 1

Q98. Which of the following is false?

(A) If $A = (1 + p/100)^t$, then $p = 100(A^{1/t} - 1)$

(B) $\sqrt{2x + 3} = x$ has only the solution $x = 3$

(C) $X^2 - rx - \delta(r + \delta) = 0$ has the solutions $x = -\delta$ and $x = r + \delta$

(D) $(p^{1/3} + 1) = 27$ implies $p = \pm \delta$

Q99. Which of the following functions are not homogenous of any degree?

(A) $5(x+y)^5 + 5$

(B) $e^{\frac{x^2}{y^2}}$

(C) $\frac{x+y}{x^2+y^2}$

(D) $3x^2y - y^3$

Q100. If $4^{40} + 4^{40} = x$, then x is:

A. 82

B. 80

C. 81

D. 160

